

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PA 97-004
reference 2

PATENT ABSTRACTS OF JAPAN

(11)Publication number : 02-290554

(43)Date of publication of application : 30.11.1990

(51)Int.Cl.

G01N 33/535

(21)Application number : 01-233999

(71)Applicant : MEIDENSHA CORP

(22)Date of filing :

08.09.1989

(72)Inventor : YOKOYAMA KAZUE

AOYANAGI SHIGEO

MATSUYUKI AKIRA

KUSUMI MIYOKO

(30)Priority

Priority number : 64 30237

Priority date : 09.02.1989

Priority country : JP

(54) REAGENT FOR ENZYME IMMUNOASSAY AND PRODUCTION THEREOF

(57)Abstract:

PURPOSE: To improve the sensitivity and reproducibility of the measuring method by forming a labeling enzyme of glucose oxidase GOD and constituting the reagent of enzyme labeled antibodies formed by combining 3 molecules of an antibody or antigen to one molecule of the GOD.

CONSTITUTION: This reagent is prepd. by mixing the GOD and the antibodies at a ratio of two-fold the number of the antibodies to be introduced into one molecule of the GOD and is in the form of a soln. or powder. The enzyme labeled antibodies are prepd. by mixing the GOD and the antibodies at a ratio of 1:2, 1:4, 1:6 (molar ratio) in the case of obtaining the enzyme labeled antibodies of 1, 2, 3 numbers of the antibodies,

respectively. The enzyme labeled antibodies of mainly 1, 2, 3 numbers of the antibodies are obtd. at the high yield and with the good reproducibility by preparing the above-mentioned antibodies at the above-mentioned ratio. The enzyme labeled antibodies prepd. in such a manner are separated from the unreacted antibodies (about 45,000mol.wt.) and further the unreacted GOD (about 150,000mol.wt.) and the enzyme labeled antibodies of the different numbers of the antibodies generated as a side reaction when the column for high-performance gel chromatograph having about $\geq 600,000$ expulsion threshold and about $\geq 15,000$ theoretical step number is used.

LEGAL STATUS

[Date of request for examination]

[Date of sending the examiner's decision of rejection]

[Kind of final disposal of application other than the examiner's decision of rejection or application converted registration]

[Date of final disposal for application]

[Patent number]

[Date of registration]

[Number of appeal against examiner's decision of rejection]

[Date of requesting appeal against examiner's decision of rejection]

[Date of extinction of right]